

Use of Electroactivated Super-Oxidized Water (HYDROCYN aqua®) as a wound cleanser in the treatment of hard-to-heal wounds: a case series

Abstract: The aim of this prospective case series is to examine the efficacy of electroactivated super-oxidized water (EASW) in wound healing. Five patient case studies treating a range of wound with EASW (Brand: HYDROCYN aqua) were selected to evaluate the efficacy. The wound healing process was documented by photographing the wound and measuring its size. The wound size was reduced, and granulation tissues formed in all patients after 2 to 3 weeks of EASW use. All of the cases demonstrated satisfactory results, with favourable clinical outcomes.

Key words:

- Antimicrobial
- Chronic wound
- Electroactivated
- Super-oxidized water
- Wound healing

Harikrishna KR Nair, Professor and Head of the Wound Care Unit, Department of Internal Medicine, Hospital Kuala Lumpur Malaysia;

Ranjeni Krishnen, PhD, Product Development Manager, Bactiguard South East Asia Sdn. Bhd., Penang, Malaysia;

Nor Azimah bt Wahab, Wound Care Unit, Segamat Hospital, Malaysia; **Chew Shao Yann**, MBBS, head of unit, wound care unit, Wound Care Unit, Melaka Hospital, Malaysia;

Farah Hana Hasan, MBBCh, Medical Officer, Head of Wound care unit, Kajang Hospital, Malaysia; **Nilrusha**, MBBS, Medical Officer, Wound Care Unit, Hospital Keningau, Sabah, Malaysia

Hard-to-heal wounds are characterised as wounds that do not heal in an orderly and timely manner with normal care (Troxler et al, 2006; Milne et al, 2020). Infection is the most common reason for hard-to-heal wounds stalling during the inflammatory phase, and it has a significant social and economic impact, lowering the quality of life for millions of patients globally. (Tenke et al, 2017; Zimmerli and Sendi, 2017). The possibility of a wound becoming infected is high since the wound surface is densely populated with a range of bacterial species (Bowler et al, 2001). Furthermore, the presence of fluid and nutrients in wound exudate and the moist condition, creates an excellent ecosystem for bacterial growth. Therefore, if the goal of therapy is to prevent wound infection and the resulting rise in patient morbidity, keeping the wound bacterial load at a threshold at which the host maintains control is crucial. Accordingly, a quality wound care plan is a fundamental prerequisite for wound healing and effective infection control in acute and chronic wounds (Atiyeh et al, 2009).

A quality wound care regimen begins with a good wound bed preparation with the goal of achieving an ideal wound-healing environment to either expedite endogenous healing or promote the effectiveness of various therapeutic procedures (Wolcott and Fletcher, 2008). A study by Schultz et al

(2003) highlighted that tissue management, inflammation and infection control, moisture balance, and epithelial (edge) advancement are the four parts of wound bed preparation that must be dealt in a systematic manner in order for wounds to heal. In accordance with this, wound cleansing can aid in the achievement of wound bed preparation goals by assisting in the removal of loose particles to establish ideal local conditions for wound healing by eliminating exudate and other debris. Irrigation is the recommended method of wound cleansing because it may remove debris and bacteria from the site while minimising stress to the wound bed. Normal sterile saline and water have traditionally been used in wound cleansing since they are nontoxic and do not disrupt healing tissues. However, in the absence of an antimicrobial component, both saline and water have no beneficial effect on wound healing because the infection remains on the wound site.

Some studies have supported the use of Electroactivated Super-Oxidized Water (EASW), also known as super-oxidized solution (SOS), as a wound cleansing agent to treat hard-to-heal wounds. There is a growing body of literature that recognises role of EASW in combating wound infection that facilitates the wound healing process (Martínez-De Jesús et al, 2007; de Angelis et al, 2013; Velazquez-Meza et al, 2015; Eftekharizadeh et al, 2016; Prabhakar et al, 2016; Singal et al, 2016; Del Rosso, and

Table 1. Inclusion and exclusion criteria	
Inclusion criteria	Patient from government hospitals who received wound care treatment/procedures with HYDROCYN aqua
	Patient is 0 to 80 years of age
	Patient meets at least one of the following indications: pressure ulcers, surgical wounds, burns (1st–2nd degree), leg ulcers
	Patient is willing and able to cooperate in the required treatment
	Patient is willing and able to complete scheduled follow-up evaluations
Exclusion criteria	Patient is unwilling or unable to give consent or to comply with the follow-up programme

Bhatia, 2018; Severing et al, 2019; Gold et al, 2020). This study discusses a case series of patients with chronic wounds who were treated with EASW (brand name: HYDROCYN aqua) to control the infection, improve granulation and accelerate healing.

Methodology

Patients case studies with various types of chronic wounds were analysed. Cases of patients treated at government hospitals throughout Malaysia were collected according to the inclusion and exclusion criteria described in [Table 1](#).

Electro-activated superoxidised water (brand name: HYDROCYN aqua®, Bactiguard SEA Sdn. Bhd.) was used to cleanse the wound before proceeding with an appropriate secondary dressing based on the wound condition.

The wound was assessed and managed with every other day (EOD) or daily dressing with EASW based on the wound condition. Firstly, gauze soaked in EASW was applied to the wound for 10 to 15 minutes to remove microorganisms and to perform autolytic debridement to soften slough. The bacteriostatic hydrogel was then applied evenly on the entire wound surface for those wounds that needed moisture. The decision on applying hydrogel was based

on a combination of clinicians' experience and preference in treating such wounds. The wound healing progress was recorded by capturing wound pictures using digital photography and measuring wound size using a paper ruler.

Results

A total of five wounds were included in this case series and there were three males and two females patients. The wounds' aetiologies included surgical wounds, venous leg ulcers, diabetic foot ulcers, and traumatic wounds ([Table 2](#)). Of the five cases, four used a sodium hypochlorite-based hydrogel (brand name: HYDROCYN aqua gel) after cleaning with EASW. Each patients' demographic data such as age, gender and wound type were recorded. [Table 2](#) illustrated the basic patient demographic. All the wounds (n=5) had achieved either complete closure or improvements from the initial baseline status.

Case 1

A 60-year-old female presented with a sloughy wound with moderate exudate. HDYROCYN aqua solution was used to treat the wound, which was the result of infection after coronary artery bypass graft (CABG) at the donor site. At presentation the wound condition

Table 1. Patient demographic, wound diagnosis and treatment outcomes					
Case number	Age (years)	Gender	Wound type	Wound size	Outcome
1	60	Female	Post-CABG* donor site wound breakdown	5 cm x 3.5 cm	Healed at 10 weeks
2	66	Female	Diabetic foot ulcer: left toe ray amputation	10 cm x 5 cm	A 98% reduction in wound size at 28 weeks
3	56	Male	Diabetic foot ulcer: right 4th finger ray amputation	3 cm x 2 cm x 5 cm	Improvements in granulation tissue a three weeks
4	45	Male	Chronic venous ulcer	10 cm x 3 cm	Sign of wound healing at 12 weeks
5	58	Male	Laparotomy dehiscence	11 cm x 6 cm	Wound re-epithelisation was almost complete at 16 weeks

*CABG: coronary artery bypass graft

Case 1. Post coronary artery bypass graft (CABG) donor site wound breakdown

- A 60-year-old female, presented with a sloughy wound with moderate exudate (*Figure 1A*)
- Treatment: HYDROCYN aqua and HYDROCYN aqua Gel, every other day
- Complete re-epithelialisation seen (*Figure 1B*)

Initial condition: Harvest site infection after CABG procedure
size of wound: 5 cm x 3.5 cm



After treatment: Wound was shrinking well and healthy epithelisation was observed. No exudate, clear from any sign of infection



Case 2. Diabetic foot ulcer after ray amputation

- A 66-year-old female, presented with a diabetic foot ulcer (DFU) following a left toe ray amputation. The wound was pale in colour, medium amount of exudate and had fetid odour (*Figure 2A*)
- Treatment: HYDROCYN aqua with daily dressing changes
- At week 28, the wound had almost completely healed and was 98% of the original size (*Figure 2C*).

Initial presentation: Wound appeared pale with medium exudate and odour
Size: 10cm x 5cm



After 24 weeks treatment: No exudate or odour, granulation tissue present
Size: 3cm x 2cm



After 28 weeks treatment: Wound was healing well
Size: 1cm x 1cm



appears to be sloughy with moderately exuding. HYDROCYN aqua gel was used after cleaning the wound with EASW. After 10 days of treatment, the wound had no signs of infection and or exudate, with signs of re-epithelialisation. Within a few weeks of wound management using HDYROCYN aqua solution, the wound condition improved and had almost completely healed by 10 weeks.

Case 2

A 66-year-old female, presented with a diabetic foot ulcer (DFU) following a left toe ray amputation. The wound was pale in colour, with medium amount of exudate and has fetid odour. After 24 weeks of treatment

with HYDROCYN aqua, the wound showed improvement with no exudate or odour. There were signs of healthy granulation tissue and a reduction in wound size by 70%. On week 28, the wound showed good healing progress with a reduction in wound size by 98%.

Case 3

A 60-year-old female, presented with a sloughy wound with moderate exudate. The patient had a right 4th finger amputation wound which was treated, for three weeks, using HYDROCYN aqua and HYDROCYN aqua gel. The wound was initially sloughy with mild amount of exudate, and showed an improvement after three weeks of treatment. Granulation tissue formed and the wound healed well.

Case 3. Diabetic foot ulcer: right 4th finger ray amputation

- A 56-year-old male, presented with a sloughy wound with moderate exudate (*Figure 3A*)
- Treatment: HYDROCYN aqua and HYDROCYN aqua Gel, every other day for three weeks
- After 3 weeks treatment, the wound showed improvement and granulation tissue could be seen (*Figure 3B*)

Initial condition: Slight slough, mild exudate, no odour.
Size of wound: 3cm x 2cm x 5cm (depth)



After 3 weeks treatment: Granulation formed and wound was healing well.
Size of wound: 1cm x 1cm



Case 4. Chronic venous leg ulcer

- A 45-year-old male, presented with a sloughy, exuding wound with a fetid odour (*Figure 4A*)
- Treatment: HYDROCYN aqua and HYDROCYN aqua Gel, every other day
- After 12 weeks, the wound showed sign of healing with no exudate and no odour, and the wound size had reduced (*Figure 4B*).

Initial condition: Sloughy, exuding with odour
Size of wound: 10cm x 3cm



After 12 weeks treatment: No exudate and odour, and wound size had reduced and was healing well
Size of wound: 3cm x 1cm



Case 4

A 45-year-old male, presented with a sloughy, exuding wound with a fetid odour. HYDROCYN aqua and gel was used to manage the chronic venous leg ulcer. After 12 weeks of application, the wound showed sign of healing with no exudate and no odour. The wound size had also reduced.

Case 5

A 58-year-old male presented with a laparotomy dehiscence wound. HYDROCYN aqua in combination of HYDROCYN aqua gel was used every other day. The wound site had no exudate and was odour less. After 16 weeks of application, wound re-epithelisation was almost complete.

Discussion

Most wounds should heal within four weeks (depending on their size and the patient's underlying comorbidities), but chronic hard-to-heal wounds may take longer or may not heal at all if proper treatment is not provided (Vowden, 2011). The intricacies of a wound, such as moisture control, the existence of tunnelling/undermining, and/or the prevalence of infection, particularly biofilm, make hard-to-heal wounds a struggle for health professionals to treat. In fact, if not treated properly these wounds may precede catastrophic outcomes such as limb amputations or even death.

Commonly, health professionals use wound bed preparation with the TIME tool to evaluate

Case 5. Laparotomy dehiscence

- A 58-year-old male, presented with a laparotomy dehiscence wound (Figure 5A)
- Treatment: HYDROCYN aqua and HYDROCYN aqua Gel, every other day
- At 16 weeks, re-epithelialisation was observed. (Figure 5B)

Initial condition: Dehiscence wound, no exudate, no odour
Size of wound: 11cm x 6cm



After 16 weeks: Re-epithelialisation was seen and the wound was healing well. Size of wound: 4cm x 2cm



and manage chronic wounds. It equips the treating health professional with a methodical way to treat chronic wounds, guaranteeing that reasonable therapies are applied, and making sure that responses are tracked and treated appropriately (Harries et al, 2016). Halim et al (2012) state that to improve a patient's capacity for healing, systemic factors must be treated before the wound bed is prepared. The initial phase of wound bed preparation is tissue management, with the objective of restoring a viable wound base with a functional extracellular matrix. In this phase, wound cleanser solution plays an important role because it is used to debride (remove) necrotic tissue, bacteria, and cells that are in the way of the healing process. This helps prevent infection and tissue loss.

In this study, each of the five cases showed decreased slough, exudate, wound size and the growth of healthy granulation tissue with new epithelialisation. EASW facilitates the mechanical removal of dead, damaged, and contaminated tissues from the wound and its surrounding area, where the microorganisms are rinsed away with solution. In addition to its primary mechanical cleansing action, the solution protects the wound from microbial invasion by reducing the microbial load at the wound upon contact with the solution. EASW contains the well-known antimicrobial agents hypochlorite ion (HOCl) and reactive oxygen species (ROS), and it kills microorganisms by denatured bacterial cell wall proteins (Kapur and Marwaha, 2011). EASW generates an unbalanced osmolarity that is detrimental to single-celled organisms due to the osmolarity

discrepancy between the concentrations of the ions in the solution and the concentrations of the same ions in the bacterial cell. Since multicellular organisms are not susceptible to such osmolarity fluctuations, the solution is non-cytotoxic. Once the membrane of a single cell is compromised, the ions in the solution also denature the bacterial proteins (Banwart, 1989; Armstrong et al, 2015). These attributes of EASW help to foster conditions that are favourable for the body's natural process of wound healing to take place.

Furthermore, usage of sodium hypochlorite-based hydrogel (HYDROCYN aqua gel) once cleansed with EASW has demonstrated promising results in the treatment of hard-to-heal wounds. These observations are consistent with those of previous research, indicating that the use of hydrogel creates a moist wound environment that promotes chronic wound healing (Singh et al, 2013; Wu et al, 2017; Gupta et al, 2019). The hydrogel creates a moist wound environment to promote wound healing by preventing dehydration and accelerating keratinocyte proliferation and migration, as well as softening hard, dry eschar (Sharman, 2003). Notably, the presence of sodium hypochlorite as an active ingredient in the hydrogel provides a bacteriostatic effect that prevents infection. To date, several studies have confirmed the antimicrobial efficacy of sodium hypochlorite (Antonelli et al, 2019; Ballal et al, 2019; Severing et al, 2019; Vajrabhaya et al, 2022).

Limitation

This study lacks a control or active comparison, patient and physician blinding, and

standardised procedures. In addition, this report is based solely on a small sample size of five case series. Therefore, it is suggested to conduct a randomised clinical trial to investigate the wound healing rate with increased sample size.

Conclusion

The use of EASW (HYDROCYN aqua) as a wound cleanser improved the wound healing process in every case in this study. Moreover, application of the bacteriostatic HYDROCYN aqua gel after wound bed preparation served as a barrier to infection and assisted in maintaining a clean wound bed. This retrospective case series demonstrates the efficacy of EASW as a standard treatment for complex chronic wounds, highlighting positive healing outcomes. The encouraging results emphasise the need for a randomised clinical trial to evaluate the efficacy of EASW in accelerating chronic wound healing.

WAS

Declaration of interest

RK is an employee of Bactiguard South East Asia Sdn.Bhd. HYDROCYN aqua and HYDROCYN aqua gel were provided by Bactiguard South East Asia Sdn.Bhd.

Acknowledgements

The authors acknowledge Bactiguard South East Asia Sdn.Bhd. for their support in this case series and for their assistance in the preparation of this manuscript.

References

- Antonelli A, Giovannini L, Baccani I et al (2019) In vitro antimicrobial activity of the decontaminant HybenX® compared to chlorhexidine and sodium hypochlorite against common bacterial and yeast pathogens. *Antibiotics (Basel)* 8(4):188. <https://doi.org/10.3390/antibiotics8040188>
- Armstrong DG, Bohn G, Glat P et al (2015) Expert recommendations for the use of hypochlorous solution. Science and Clinical Application. *Ostomy Wound Manage* 61(5):S2–19
- Atiyeh BS, Dibo SA, Hayek SN (2009) Wound cleansing, topical antiseptics and wound healing. *Int Wound J* 6(6):420–30. <https://doi.org/10.1111/j.1742-481x.2009.00639.x>
- Ballal NV, Gandhi P, Shenoy PA et al (2019) Safety assessment of an etidronate in a sodium hypochlorite solution: randomized double-blind trial. *Int Endod J* 52(9):1274–82. <https://doi.org/10.1111/iej.13129>
- Banwart GJ (1989) Control of microorganisms by retarding growth. In: *Basic Food Microbiology*. Springer pp545–50.
- Bowler PG, Duerden BI, Armstrong DG (2001) Wound microbiology and associated approaches to wound management. *Clin Microbiol Rev* 14(2):244–69. <https://doi.org/10.1128%2FCMR.14.2.244-269.2001>
- de Angelis B, Lucarini L, Agovino A et al (2013) Combined use of super-oxidised solution with negative pressure for the treatment of pressure ulcers: case report. *Int Wound J* 10(3):336–9. <https://doi.org/10.1111/j.1742-481x.2012.00982.x>
- Del Rosso JQ, Bhatia N (2018) Status report on topical hypochlorous acid: clinical relevance of specific formulations, potential modes of action, and study outcomes. *J Clin Aesthet Dermatol* 11(11):36–39
- Eftekhariadeh F, Dehnavieh R, Hekmat SN, Mehrolhassani MH (2016) Health technology assessment on super oxidized water for treatment of chronic wounds. *Med J Islam Repub Iran* 30:384
- Gold MH, Andriessen A, Bhatia AC et al (2020) Topical stabilized hypochlorous acid: The future gold standard for wound care and scar management in dermatologic and plastic surgery procedures. *J Cosmet Dermatol* 19(2):270–7.
- Gupta A, Kowalczyk M, Heaselgrave W et al (2019) The production and application of hydrogels for wound management: A review. *European Polymer Journal* 111:134–151. <https://doi.org/10.1016/j.eurpolymj.2018.12.019>
- Halim AS, Khoo TL, Saad AM (2012) Wound bed preparation from a clinical perspective. *Indian J Plast Surg* 45(2):193–202. <https://doi.org/10.4103%2F0970-0358.101277>
- Harries RL, Bosanquet DC, Harding KG (2016) Wound bed preparation: TIME for an update. *Int Wound J* 13(S3):8–14. <https://doi.org/10.4103%2F0970-0358.101277>
- Kapur V, Marwaha AK (2011) Evaluation of effect and comparison of superoxidised solution (oxum) v/s povidone iodine (betadine). *Indian J Surg* 73(1):48–53. <https://doi.org/10.1007/s12262-010-0189-y>
- Martínez-De Jesús FR, Ramos-De la Medina A, Remes-Troche JM et al (2007) Efficacy and safety of neutral pH superoxidised solution in severe diabetic foot infections. *Int Wound J* 4(4):353–62. <https://doi.org/10.1111/j.1742-481x.2007.00363.x>
- Milne J, Searle R, Styche T (2020) The characteristics and impact of hard-to-heal wounds: results of a standardised survey. *J Wound Care* 29(5):282–8. <https://doi.org/10.12968/jowc.2020.29.5.282>
- Prabhakar KBS, Purushotham G, Uma K (2016) Comparison of super-oxidized solution versus povidine iodine in management of infected diabetic ulcers: our experience. *IAIM* 3(5):151–8
- Schultz GS, Sibbald RG, Falanga V et al (2003) Wound bed preparation: a systematic approach to wound management. *Wound Repair Regen* 11:51–28. <https://doi.org/10.1046/j.1524-475x.11.s2.1.x>
- Severing AL, Rembe JD, Koester V, Stuermer EK (2019) Safety and efficacy profiles of different commercial sodium hypochlorite/hypochlorous acid solutions (NaClO/HClO): antimicrobial efficacy, cytotoxic impact and physicochemical parameters in vitro. *J Antimicrob Chemother* 74(2):365–72. <https://doi.org/10.1093/jac/dky432>
- Sharman D (2003) Moist wound healing: a review of evidence, application and outcome. *The Diabetic Foot* 6(3):112–20. <https://tinyurl.com/5x22wmru> (accessed 12 February)
- Singal R, Siddharth DHAR, Zaman M et al (2016) Comparative evaluation of intra-operative peritoneal lavage with super oxidized solution and normal saline in peritonitis cases; randomized controlled trial. *Maedica (Bucur)* 11(4):277–85
- Singh B, Sharma S, Dhiman A (2013) Design of antibiotic containing hydrogel wound dressings: biomedical properties and histological study of wound healing. *Int J Pharm* 457(1):82–91. <https://doi.org/10.1016/j.>

ijpharm.2013.09.028

Tenke P, Mezei T, Bóde I et al (2017) Catheter-associated urinary tract infections. *Eur Urol Suppl* 16(4):138–43. <https://doi.org/10.1016/j.eursup.2016.10.001>

Troxler M, Vowden K, Vowden P (2006) Integrating adjunctive therapy into practice: the importance of recognising 'hard-to-heal'wounds. *World Wide Wounds* 32(1):99–105. <https://tinyurl.com/58zpjfeb> (accessed 12 February)

Vajrabhaya LO, Korsuwannawong S, Ruangsawasdi N et al (2022) The efficiency of natural wound healing and bacterial biofilm inhibition of Aloe vera and Sodium Chloride toothpaste preparation. *BMC Complement Med Ther* 22(1):1–10. <https://doi.org/10.1186/s12906-022-03548-7>

Velazquez-Meza ME, Hernández-Salgado M, Sánchez-Alemán MA (2015) Evaluation of the antimicrobial

activity of a super oxidized solution in clinical isolates. *Microbial Drug Resistance* 21(4):367–72. <https://doi.org/10.1089/mdr.2014.0266>

Vowden P (2011) Hard-to-heal wounds Made Easy. *Wounds International* 2(4): <https://tinyurl.com/2s3rzjms> (accessed 12 February)

Wolcott R, Fletcher J (2008) The role of wound cleansing in the management of wounds. *Inflammation* 17(8):333–41. <https://tinyurl.com/59m4u8sp> (accessed 12 February)

Wu S, Deng L, Hsia H et al (2017) Evaluation of gelatin-hyaluronic acid composite hydrogels for accelerating wound healing. *J Biomater Appl* 31(10):1380–90. <https://doi.org/10.1177/0885328217702526>

Zimmerli W, Sendi P (2017) Orthopaedic biofilm infections. *J Pathol Microbiol Immunol* 125(4):353–64. <https://doi.org/10.1111/apm.12687>

